



2021
2022

PRODUCTS CATALOG

addvet.net



WELCOME TO CATALOG - 2021 2022



addvet.net

COMPANY PROFILE

Who we are

Established in 2014 with capital of one million dollars, ADDVET is a Jordanian company specialized in manufacturing veterinary drugs.

What we offer


A variety of veterinary medicines Powders & Liquids orals, including Antibiotics, Vitamins, Amino Acids, Anticoccidial drugs & Premixes.

Our Mission

A variety of veterinary medicines Powders & Liquids orals, including Antibiotics, Vitamins, Amino Acids, Anticoccidial drugs & Premixes.

Our Vision

To improve the world feed and food business and be part of the next generation global feed chain.



addvet.net



SUPPLEMENT

AD3E SUPER
AD3E+C
AD3E+CK
ADD-C100
ADDCHOLIN+
ADD-E120
ADD-K3100
ADDLYTE
ADDPower
ADDVITA+
ADD-VITAB+
AMINOVITA
CALPHOS+
LIVERVET
MOULDOUT



addvet.net

ADDvet
Animal Health

2

3

FOR VETERINARY USE ONLY.

AD3E SUPER

Promotion of growth rates, improvement of feed efficiency and strengthens performance.

Composition

Each 1 Litre contains:

Vitamin A	100 000 000 I.U.
Vitamin D3	20 000 000 I.U.
Vitamin E	20 000 mg

Indications

- Prevention of vitamins insufficiencies in poultry.
- Strengthen of resistance to the diseases.
- Improvement of egg production and hatchability.
- Promotion of growth rates and improves feed efficiency.

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.

1L



SUPPLEMENT

AD3E+C

Promotion of growth rates, improvement of feed efficiency and strengthens immunity.

Composition

Each 1 Litre contains:

Vitamin A	10,000,000 I.U.
Vitamin D3	2,000,000 I.U.
Vitamin E	10,000 mg
Vitamin C	20,000 mg

Indications

- Prevention and control of stress conditions.
- Prevention of vitamins insufficiency in poultry.
- Strengthen of resistance to the diseases.
- Improvement of egg production and hatchability.
- Promotion of growth rates and improves feed efficiency.

Dosage and Administration

200-250 mL per 1000 Litres of drinking water for 3-5 days or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

4

AD3E+CK

Growth rates promotion, improvement of feed efficiency and strengthens animal performance.

Composition

Each 1 Litre contains:

Vitamin A	25 000 000 I.U.
Vitamin D3	2 000 000 I.U.
Vitamin E	10 000 mg
Vitamin C	20,000 mg
Vitamin K3	20,000 mg
Distillate water up to	1000 mL

Indications

- Prevention of vitamins insufficiencies in poultry.
- Strengthen of resistance to the diseases.
- Improvement of egg production and hatchability.
- Promotion of growth rates and improves feed efficiency.
- Improvement of immune system and against heat stress.
- Plays a a role in blood clotting mechanisms.

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

5

ADD-C100

Reliable for stress condition of high temperature and better immune system function.

Composition

Each 1 Litre contains:
 Vitamin C 100,000 mg
 Distillate water up to 1000 mL

Vitamin C is vital for proper immune system activity as it acts as an enzymatic cofactor for eight different. It also has antioxidant activity for proper operation of several physiological functions.

Indications

- For heat stress cases caused by extreme high temperature.
- For better immune system against various diseases.
- Improves feed intake, weight gain, and egg production.

Dosage and Administration

1000 mL per 2000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
 STORE PRODUCT IN A COOL PLACE.

6

ADDCHOLIN+

Enhance and improve the liver function of the poultry and preventing perosis.

Composition

Each 1 Litre contains:
 Choline chloride 22 000 mg
 L-Carnitine 20 000 mg
 Betaine 80 000 mg
 Magnesium sulfate 50 000 mg

Indications

- Prevention of perosis and liver enlargement in chicks, helps in utilization and outward movement of fat from the liver and prevent from condition known as fatty liver.
- Growth promotion and strengthening the immune system.
- Improves production performance and increases intracellular water retention.
- Magnesium sulfate increases the water level in the intestine.

Dosage and Administration

500 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

7

ADD-E120

Powerful antioxidant for crazy chick disease, exudative diathesis and muscular dystrophy.

Composition

Each 1 Litre contains:

Vitamin E	120.000 mg
Sodium selenite	60 mg
Distillate water up to	1000 mL

Indications

- Vitamin E, an antioxidant for normal neurological functions.
- Selenium is essential for glutathione peroxidase enzyme composition that protects cells from destroying oxidising agents.
- For Vitamin E and Selenium deficiency diseases such as encephalomalacia, muscular dystrophy, and exudative diathesis.
- For stress conditions, hatchability and infertility problems.

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

8

ADD-K3100

A blood clotting mechanism for bleeding disorders due to Vitamin K3 deficiency.

Composition

Each 1 Litre contains:

Vitamin K3	100.000 mg
Distillate water up to	1000 mL

Indications

Vitamin K3 is essential for synthesis of prothrombin, thus it plays an important role in clotting mechanisms and also has a protective effect against coccidiosis.

Deficiency of vitamin K3 may cause an increase of blood spots in eggs, haemorrhages in the legs and breast and a failure of blood clotting.

Dosage and Administration

500-1000 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

9

ADDLYTE

Important role as an electrolyte to regulate cellular functions and maintain osmotic balance.

Composition

Each 1 Litre contains:

Ammonium chloride	20.000 mg	Cobalt sulphate	3.000 mg
Magnesium sulphate	20.000 mg	Potassium iodite	1.000 mg
Copper sulphate	10.000 mg	Sodium selenite	50 mg
Zinc sulphate	15.000 mg	Vitamin C	5.000 mg
Iron sulphate	5.000 mg		

Indications

- Balanced formula for dehydration and electrolytes imbalance.
- For stress situations such as diarrhea and high temperature.
- Provides osmotic balance and detoxification in body.

Dosage and Administration

500-1000 mL per 1000 Litres of drinking water for 3-5 days or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

10

11

FOR VETERINARY USE ONLY.

ADDPOWER

Functions as an energy activation nutrient to assist tired lethargic animals and dehydration.

Composition

Each 1 Litre contains:

Dextrose	400.000 mg
Vitamin B1	5.000 mg
Vitamin B2	200 mg
Vitamin B6	2.000 mg
Vitamin B12	5 mcg
Distillate water up to	1000 mL

Indications

- Maintenance of fluid balance by adjusting the water intake.
- Replace lost fluids and provide carbohydrates to the body.
- Regulation of acid-base balance by maintaining blood pH.
- Helps in perosis cases and animals in lactation.
- Helps in stress cases and hatchability problems.

Dosage and Administration

500 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

ADDVITA+

Combination of well-balanced formula for vitamins and amino acids deficiencies in animals.

Composition

Each 1 Litre contains:

Vitamin A	15 000 000 I.U.	D-Panthenol	2 820 mg
Vitamin D3	1 000 000 I.U.	Choline Chloride	70 000 mg
Vitamin E	6 000 mg	Betaine	3 000 mg
Vitamin B1	1 500 mg	Lysine	6 000 mg
Vitamin B2	1 660 mg	Methionine	3 000 mg
Vitamin B6	1 000 mg	Threonine	1 500 mg
Vitamin B12	10 mg	Tryptophan	250 mg
Niacin	15 000 mg	Sorbitol	50 000 mg
Biotin	900 mg	Sodium propionate	4 200 mg
Folic acid	200 mg	Purified water up to	1 Litre

Indications

- Formulation for improved growth rate, feed utilization and egg production.
- Helps in vitamins and amino acids deficiencies that causes bad health.
- Improve resistance to various stress conditions (caused by vaccination, diseases, transport, high humidity, high temperatures or extreme temperature changes).

Dosage and Administration

250 mL per 1000 Litres of drinking water.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

12

ADD-VITAB+

Function as a nutritional source of B-Complex vitamin deficiencies in animals.

Composition

Each 1 Litre contains:

Vitamin B1	2 000 mg	Vitamin K3	5 000 mg
Vitamin B2	3 500 mg	Vitamin B12	12 mg
Vitamin PP	22 000 mg	Folic acid	500 mg
Vitamin B5	13 000 mg	Choline chloride	20 000 mg
Vitamin B6	6 000 mg	Betaine	30 000mg

Indications

- Helps in retarded growth during antibiotic treatment.
- Improve body defences during stress conditions.
- Improve feed conversion.
- In stress conditions (caused by vaccination, diseases, transport, high humidity, high temperatures or extreme temperature changes).

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

13

AMINOVITA

Balanced formula of vitamins and amino acids for the general health welfare of poultry.

Composition

Each 1 Litre contains:

Vitamin A	15 000 000 I.U.	D-Panthenol	2 820 mg
Vitamin D3	1 000 000 I.U.	Choline Chloride	70 000 mg
Vitamin E	6 000 mg	Betaine	3 000 mg
Vitamin B1	1 500 mg	Lysine	6 000 mg
Vitamin B2	1 660 mg	Methionine	3 000 mg
Vitamin B6	1 000 mg	Threonine	1 500 mg
Vitamin B12	10 mg	Tryptophan	250 mg
Niacin	15 000 mg	Sorbitol	50 000 mg
Biotin	900 mg	Sodium propionate	4 200 mg
Folic acid	200 mg	Purified water up to	1 Litre

Indications

- Prevents and treats various vitamins and amino acids deficiencies.
- Achieves faster growth rate and improves weight gain plus feed conversion.
- Helps in conditions of stress like heat temperature or diseases.
- Stimulates the metabolism and increases egg productivity.

Dosage and Administration

250 mL per 1000 Litres of drinking water.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

14

CALPHOS+

Function to develop stronger animal bones development and increase milk production.

Composition

Each 1 Litre contains:

Phosphorus	110 000 mg	Zinc	5 000 mg
Magnesium	22 000 mg	Copper	200 mg
Calcium	28 000 mg		

Indications

- Maintain good shell quality and strong bones development.
- High levels of phosphorus in the blood will inhibit the mobilization of calcium from bones.
- Prevent calcium deficiency in laying hens results in reduced shell quality and subsequently cause osteoporosis.
- Increase milk production and faster growth.

Dosage and Administration

500 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

15

LIVERVET

Effective in fatty liver syndrome and improves the animals' overall liver function.

Composition

Each 1 Litre contains:

Dextrose	100.000 mg	Niacine	30.000 mg
Vitamin B1	200 mg	Methionine	20.000 mg
Vitamin B6	200 mg	Vitamin E	20.000 mg
Vitamin B12	30 mcg	Vitamin C	10.000 mg
Choline chloride	70.000 mg	Distillate water up to	1000 mL

Indications

- Prevents fatty liver syndrome and improves liver functions.
- Improves carbohydrate metabolism for proper functioning of nervous tissue, cardiac muscle and enzyme activities.
- Important role in metabolism, growth rate and feed utilization.
- Stimulates regeneration of hepatic cells and better productivity.

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

16

MOULDOUT

Binds mycotoxins, removing them from the animal's digestive tract and improve growth.

Composition

Each 1 Litre contains:

Yeast extracts	200 000 mg	Citric acid	70 000 mg
Phosphoric acid	70 000 mg	Betaine	80 000 mg
Lactic acid	70 000 mg		

Indications

- Suppression of proliferation of intestinal harmful bacteria by reducing pH in gastrointestinal tract.
- Growth promotion, better weight gain, and improvement of feed efficiency.
- Prevention of dietary diarrhea by promoting digestion.
- Improve feed consumption and feed conversion rate.

Dosage and Administration

250 mL per 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

17

ANTIBIOTIC

ADDCIPRO
ADDENRO
ADDFLOR
ADDFOS
ADDTILMO
ENROHEXIN
TYFOS
TYLOHEXIN
ADDAMOX
ADDDOX
ADDOXY
ADDTHRO

AMOXYCOL
AMPICOL
DOXYCOL
GENTDOXY
GENTTYLO
TYLODOXY
TYLOTHRO
ADDAMPI-50
ADDCOL12
TYLOFORT



addvet.net

ADDvet
Animal Health

ADDCIPRO

Acts bactericidally to treat gastrointestinal, respiratory and urinary tract infections.

Composition

Each 1 mL contains:
Ciprofloxacin 100 mg

Ciprofloxacin belongs to the group of quinolones and acts bactericidally against mainly Gramnegative bacteria like Campylobacter, E. coli, Haemophilus, Mycoplasma, Pasteurella and Salmonella spp.

Indications

To treat ciprofloxacin sensitive microorganisms in poultry, calves, goat and sheep that cause gastrointestinal, respiratory and urinary tract infections.

Dosage and Administration

1 Litre / 1000 Litres of drinking water or as advised by a veterinarian.



1L

ADDENRO

Acts bactericidally to treat gastrointestinal, respiratory and urinary tract infections.

Composition

Each 1 mL contains:
Enrofloxacin 200 mg

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp.

Indications

To treat enrofloxacin sensitive microorganisms in poultry, calves, goat and sheep that cause gastrointestinal, respiratory and urinary tract infections.

Dosage and Administration

500 mL / 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

20

ADDFLOR

A bacteriostatic to treat gastrointestinal, and respiratory tract infections in animals.

Composition

Each 1 mL contains:
Florfenicol 100 mg

Florfenicol is a synthetic broad-spectrum antibiotic that is effective against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol, a fluorinated derivative of chloramphenicol, acts by inhibiting protein synthesis at the ribosomal level and is bacteriostatic.

Indications

Preventive and therapeutic treatment of gastrointestinal and respiratory tract infections, caused by florfenicol sensitive microorganisms such as Actinobaccillus spp., Pasteurella spp., Salmonella spp. and Streptococcus spp. in poultry.

Dosage and Administration

1 Litre / 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

21

ADDFOS

Broad-spectrum bactericidal antimicrobial of fosfonic acid family to treat diseases.

Composition

Each 1 mL contains:
Fosfomycin (as sodium) 100 mg

Fosfomycin is a broad spectrum bactericidal antimicrobial belonging to the fosfonic acid family, recommended as an aid for the treatment of bacterial diseases of poultry.

Indications

For the treatment of bacterial infections of poultry susceptible to the action of fosfomycin such as Infectious coryza, fowl cholera, colibacillosis, ORT, CRD, staphylococcal infections, fowl typhoid, and salmonellosis.

Dosage and Administration

500 mL / 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

22

ADDTILMO

Broad-spectrum semi-synthetic bactericidal macrolide antibiotic to treat respiratory diseases in poultry.

Composition

Each 1 mL contains:
Tilmicosin phosphate 250 mg

Tilmicosin is a macrolide antibiotic. It is used in veterinary medicine for the treatment of bovine respiratory disease and bovine respiratory disease associated with Mannheimia (Pasteurella) haemolytica.

Indications

For the control and treatment of respiratory infections associated with tilmicosin-susceptible micro-organisms such as Mycoplasma spp. Pasteurella multocida, Actinobacillus pleuropneumoniae, Actinomyces pyogenes and Mannheimia haemolytica in chickens and turkeys.

Dosage and Administration

250 mL / 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

23

ENROHEXIN

Quinolone antibiotic with added bromhexine to treat respiratory tract diseases in animals.

Composition

Each 1 mL contains:

Enrofloxacin 100 mg Bromhexine 10 mg

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella and Mycoplasma spp. Bromhexine is a mucolytic.

Indications

For the treatment and control of respiratory infections associated with Mycoplasma spp., Pasteurella multocida., Haemophilus paragalinarum, Chronic Respiratory Disease (CRD), in some infections produced by susceptible strains of Staphylococcus, and also for other enrofloxacin-susceptible bacteria.

Dosage and Administration

1 Litre / 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

24

TYFOS

Bacteriostatic action against mainly Gram-positive and Gram-negative bacteria.

Composition

Each 1 mL contains:

Tylosin 200 mg
Fosfomycin 100 mg

Indications

It is a combination of broad-spectrum antibiotic, indicated for the treatment of poultry diseases, caused by Gram-positive, Gram-negative and Mycoplasmas. Indicated in cases of: mycoplasmosis, colibacillosis, chronic respiratory disease in poultry, Staphylococcus infections, Listeriosis, Avian Cholera, infectious Coryza, and diseases caused by Fosfomycin and/or Tylosin susceptible organisms.

Dosage and Administration

1 Litre / 1000 Litres of drinking water or as advised by a veterinarian.



1L

FOR VETERINARY USE ONLY.

25

TYLOHEXIN

Macrolide antibiotic with added bromhexine to treat respiratory tract diseases.

Composition

Each 1 mL contains:

Tylosin 200 mg
Bromhexine 10 mg

Indications

For the treatment and control of respiratory infections associated with Mycoplasma spp., Pasteurella multocida., Haemophilus paragalinarum, Chronic Respiratory Disease (CRD), in some infections produced by susceptible strains of Staphylococcus, and also for other enrofloxacin-susceptible bacteria.

Dosage and Administration

1 Litre / 1000 Litres of drinking water or as advised by a veterinarian.



1L

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

26

ADDAMOX

Broad-spectrum penicillin with a bactericidal action against both Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Amoxicillin 300 mg

The bactericidal action is due to inhibition of cell wall synthesis. Amoxicillin is mainly excreted in urine. A major part can also be excreted in bile.

Indications

Gastrointestinal, respiratory and urinary tract infections caused by amoxicillin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500 mg

FOR VETERINARY USE ONLY.

27

ADDDOX

Acts bacteriostatic against many Gram-positive and Gram-negative bacteria in animals.

Composition

Each 1 gm contains:
Doxycycline cyclate 500 mg

Doxycycline belongs to the group of tetracyclines and acts bacteriostatic against many Gram-positive and Gram-negative bacteria. The action of doxycycline is based on inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by doxycycline sensitive micro-organisms, like Bordetella, Campylobacter, Chlamydia, E. coli, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

250 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

28

ADDOXY

Broad-spectrum tetracycline with a bactericidal action against both Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Oxytetracycline 500 mg

The mode of action of oxytetracycline is based on inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by oxytetracycline sensitive bacteria like Bordetella, Bacillus, Corynebacterium, Campylobacter, E. coli, Haemophilus, Pasteurella, Salmonella, Staphylococcus and Streptococcus spp. and Mycoplasma, Rickettsia and Chlamydia spp. in various animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

FOR VETERINARY USE ONLY.

29

ADDTHRO

Reliable for Chronic Respiratory Disease (CRD) caused by Mycoplasma infections in poultry.

Composition

Each 1 gm contains:
Erythromycin thiocyanate 350 mg

Erythromycin binds reversibly to the 50S subunit of ribosomes, resulting in the blockage of transpeptidation or translocation reactions, leading to impaired and/or inhibited protein synthesis and to inhibited cell growth.

Indications

Chronic Respiratory Disease (CRD) caused by Mycoplasma infections, and for the treatment of infectious respiratory diseases due to erythromycin-sensitive bacteria such as Staphylococcus aureus (e.g. infectious arthritis), Haemophilus paragallinarum (e.g. infectious coryza) and Pasteurella multocida (e.g. pasteurellosis; pneumonia) in poultry.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

30

AMOXYCOL

Quality duo medicinal combination with bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Amoxicillin trihydrate 200 mg
Colistin sulphate 1200000 I.U.

Indications

Gastrointestinal, respiratory and urinary tract infections caused by amoxicillin and colistin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in animals.

The action of amoxicillin is based on inhibition of cell wall synthesis. Colistin interact strongly with phospholipids and penetrate into and disrupt the structure of cell membranes.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

FOR VETERINARY USE ONLY.

31

AMPICOL

Quality duo combination with bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Ampicillin trihydrate	200 mg
Colistin sulphate	1200000 I.U.

Indications

Gastrointestinal and urinary tract infections caused by ampicillin and colistin sensitive micro-organisms, like Campylobacter, Clostridium, Corynebacterium, E. coli, Erysipelothrix, Haemophilus, Pasteurella, Salmonella, penicillinase negative Staphylococcus and Streptococcus spp. in animals.

It can also be used for the treatment of clinical bovine mastitis in the lactating cow. It is effective for diarrhea and toxicity cases.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500 mg

**KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.**

32

DOXYCOL

Quality duo medicinal combination acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Doxycycline hyclate	200 mg
Colistin sulphate	1200000 I.U.

Indications

Gastrointestinal and respiratory infections (CRD) caused by microorganisms sensitive to colistin and/or doxycycline like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

The action of doxycycline is based on inhibition of bacterial protein synthesis. Colistin interact strongly with phospholipids and penetrate into and disrupt the structure of cell membranes.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500 mg

FOR VETERINARY USE ONLY.

33

GENTDOXY

Duo antibiotic class that acts bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Doxycycline hyclate 200 mg
Gentamicin sulphate 200 mg

The action of doxycycline and gentamicin is based on inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by microorganisms sensitive to doxycycline and/or gentamicin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500 mg

**KEEP OUT OF REACH OF CHILDREN.
 STORE PRODUCT IN A COOL PLACE.**

34

GENTTYLO

Duo antibiotic class that acts bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Tylosin tartrate 200 mg
Gentamicin sulphate 200 mg

The action of both tylosin and gentamicin is based on inhibition of bacterial protein synthesis.

Indications

Gastrointestinal and respiratory infections caused by microorganisms sensitive to tylosin and/or gentamicin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500 mg

FOR VETERINARY USE ONLY.

35

TYLODOXY

Quality duo medicinal combination acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Tylosin tartrate	120 mg
Doxycycline hyclate	200 mg


The action of tylosin and doxycycline is based on inhibition of bacterial protein synthesis. They binds reversibly to the 50S subunit of ribosomes.

Indications

Gastrointestinal and respiratory infections (CRD) caused by microorganisms sensitive to tylosin and/or doxycycline like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

36

TYLOTHRO

Quality duo macrolide class acting bacteriostatic against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:

Tylosin tartrate	120 mg
Erythromycin thiocyanate	200 mg

The action of tylosin and erythromycin is based on inhibition of bacterial protein synthesis. They binds reversibly to the 50S subunit of ribosomes.

Indications

Gastrointestinal and respiratory infections (CRD) caused by microorganism sensitive to tylosin and/or erythromycin like Bordetella, Campylobacter, Chlamydia, E. coli, Klebsiella, Haemophilus, Mycoplasma, Pasteurella, Rickettsia, Salmonella, Staphylococcus and Streptococcus spp. in animals.

Dosage and Administration

500 gm / 1000 Litres of drinking water or as advised by a veterinarian.



500
mg

FOR VETERINARY USE ONLY.

37

ADDAMPI-50

Quality duo combination with bactericidal action against many Gram+ve and Gram-ve bacteria.

Composition

Each 1 gm contains:
Ampicillin (Trihydrate) 500 mg

Indications

Ampicillin is indicated for the treatment of infections caused by E.coli, Clostridium, Salmonella, Brucella, Proteus, Klebsiella, Strepto and Staphylococci, Corynebacterium, Haemophilus, Pasteurella, and Shigella in the different animals. Ampicillin 50% is recommended for uses in cases of Bacterial enteritis and Necrotic enteritis in poultry. It is also indicated Against Pneumonia, Septicemia, Diarrhea, and Scours in calves and lambs.

Dosage and Administration

100 gm / 500 Litres of drinking water or as advised by a veterinarian.



500 mg

KEEP OUT OF REACH OF CHILDREN.
STORE PRODUCT IN A COOL PLACE.

38

ADDCOL12

Acts bacteriostatic against many Gram-positive and Gram-negative bacteria in animals.

Composition

Each 1 gm contains:
Colistin sulphate 1200000 IU.

Indications

ADDCOL12 Powder is indicated for the treatment of enteric infections caused by gram-negative bacteria e.g. E. coli, Salmonella, Shigella in poultry, calves and lambs. Colistin are highly effective against strains of E. coli, Pseudomonas aeruginosa, Salmonella spp., Haemophilus spp., Shigella spp., Pasteurella spp., Brucella spp., Aerobacteraerogenes, and Bordetella bronchiseptica. Gram-positive bacteria are generally less sensitive. However, there are sensitive strains of Staphylococcus spp., Bacillus spp., Streptococcus pyogenes and Corynebacterium spp.

Dosage and Administration

Calves, goats and sheep:

Twice daily 5 g per 100 kg body weight for 5 - 7 days.

Poultry: 500 gm per 500 - 1000 litres of drinking water or 250 - 500 kg of feed for 5 - 7 days.



500 mg

FOR VETERINARY USE ONLY.

39

ANTIBIOTIC & SUPPLEMENT

addvet.net



KEEP OUT OF REACH OF CHILDREN

STORE PRODUCT IN A COOL PLACE

ADDvet
Animal Health

44

45

ADDvet
Animal Health

addvet.net



Contact

Al-Zarqa Industrial Zone [1Z4], Jordan.
info@addvet.net

